

Strategies and Methodologies of Improving Rice Varieties in Egypt

A. A. Abd Allah, Shimaa A. Badawy* and A. T. Badawi

Rice Research Department, Field Crop Research Institute, Agricultural Research Center, Gizza and *Agronomy Department, Faculty of Agriculture, Kafrelsheikh University, Kafrelsheikh, Egypt.

IN THE EXPERIMENTAL Farm of the Rice Research and Training Center (RRTC), Sakha, Kafer elshiekh, Egypt, during the period from 2005 to 2011 rice growing seasons. The present investigation was carried out to evaluate some Delta-Med materials under Egyptian conditions and develop a new rice lines using a crosses between 7 Egyptian varieties (Giza 177, Giza 178, Sakha 101, Sakha 102, Sakha103, Sakha 104 and Giza 182) and 7Spain varieties (Bala, Susan, Bahia, Bomba, Montsianejj, Senia and Tebre) to enhance the former genotype yielding ability. The results showed that there were significant differences between Egyptian varieties and Delta-Med varieties as well as among each group itself. The Egyptian varieties exceeded the Delta-Med entries in grain yield and its associated traits. For Spanish and Delta-Med genotypes, they were generally low grain yield, shorter duration and higher 1000-grain-weight. With crosses between Egyptian and Spain varieties, the former genotype yielding ability were enhanced along with a specific breeding program to be designed for Spain materials and be evaluated in Spain in the few coming years. The progenies from each cross were advanced under normal conditions with the pedigree method technique. The best selected lines from F6 generation (2010) were promoted to be grown as F7 generation in 2011 season. These lines were found to be resistant to blast disease in addition to possessing useful traits associated with high yielding ability. Phenotypic coefficient of variation (PCV) was high for all the studied character and genotypic coefficient of variability (GCV) showed the same trend as for PCV. Heritability estimates were high for all studied traits. Highly significant positive phenotypic correlation coefficients were determined between grain yield per plant and all yield components studied.

Keywords: Rice, Grain, Plant breeding, Yielding, Egypt.

Rice is the most important food crop in the world. Almost half of the world's population depends on rice as their staple food. Rice is the most important crop to directly consume by humans. With around 600 M t produced on 149 M ha in 2003, rice accounts for 23% of the world's caloric intake (Khunthasuvon *et al.*, 1998). Rice production increased by 130% between 1966 and 2000, while the population of low income countries increased by an average of 90% over the same period. The world population is predicted to reach approximately 8 billion

by 2030 and there is therefore a need to further increase rice production by 40% in the next 25 years (IRRI, 1996).

Therefore, to meet the needs of the growing world population, rice production should be increased via developing new high yielding and drought tolerant rice varieties. The rice growing area is about half million hectares with a national average yield of 10 tons/ ha in Egypt. This significantly high productivity is mainly due to the development of short duration varieties with a higher yield potential(10 t/ha), resistance to biotic and a biotic stresses, better grain quality (high milling and low amylase content) in combine with an excellent extension network through the national campaign of rice (RRTC, 2007).

As a part of the international co-operation with Delta Med countries, an initiative were carried out at the Institute of Agroambiental TerresEbre., Spain, to evaluate six of our newly released varieties, along with four checks in 2005 season. The Egyptian varieties were significantly different in their behavior under Spain conditions. The objectives of the current work are to evaluate some Spain materials under Egyptian conditions and also to start an initiative to develop new anther culture derived lines using crosses between Egyptian and Spain varieties to enhance the former genotype yielding ability. These lines, along with a specific breeding program designed for Spain materials, to be evaluated in Spain in the few coming years.

The objectives of the current work is to evaluate some Delta Med materials under Egyptian conditions and also to develop a new rice lines using a crosses between Egyptian and Spain varieties to enhance the former genotype yielding ability.

Materials and Methods

Materials

Parents

A total of 14 entries including 7 Egyptian varieties *i.e.*, Giza 177, Giza 178, Sakha 101, Sakha 102, Sakha103, Sakha 104 and Giza 182 and 7 spanish ones, namely, Bala, Susan, Bahia, Bomba, Montsianejj, Senia and Tebre.

The pedigree generations of Delta-Med materials

A pedigree breeding program specifically to the Delta Med materials have reached up to F7 stage in 2011 rice season. The summary of the breeding program with Delta med materials are presented in Table 1 and Table 2. A total of 45 F7 lines were selected according to their desirable characters such as high yielding ability, blast resistance, stem borer resistance, grain weight and grain quality characters. These lines will be promoted to be grown as yield trial experiments in 2012 rice season. One set of this F7 lines will be distributed to be grown in the different locations as a verification trials through technology transfer program and after that can be sent to the Delta Med countries for evaluation there.

Egypt. J. Agron. **34**, No.2 (2012)

TABLE 1. Summary of the pedigree generations of Delta-Med materials (2010).

Year	Generation	Planted		Selected	
		Lines	Crosses	Lines	Crosses
2006	F1	-	24	-	24
	F2	323	24	286	10
2007	F1	-	16	-	16
	F2	500	8	75	7
	F3	263	10	212	10
2008	F2	380	16	158	15
	F3	204	7	106	7
	F4	230	10	115	10
2009	F3	148	14	102	10
	F4	96	6	134	6
	F5	110	9	100	8
2010	F4	101	10	85	9
	F5	128	6	66	3
	F6	75	3	45	3
2011	F5	85	9	30	9
	F6	66	3	25	3
	F7	45	3	6	3

TABLE 2. Summary of the pedigree generations of Delta-Med materials (2011).

Generation	Planted		Selected	
	Lines	Crosses	Lines	Crosses
F5	81	9	30	9
F6	43	3	25	3
F7	15	3	6	3
Total	135	15	61	15

Methods

The present investigation was carried out at the Experimental Farm of the Rice Research and Training Center (RRTC), Sakha, Kafer elshiekh, Egypt during seven successive rice growing seasons (from 2005 to 2011) to evaluate some Delta Med materials under Egyptian conditions and also to develop a new rice lines using a crosses between Egyptian and Spain varieties to enhance the former genotype yielding ability.

The tested materials were grown in randomized complete block design (RCBD) with three replications, each plot were 7 rows 5 m long with a plant spacing 20×20 cm, the nurseries were raised till age of 30 days, then, they were transplanted with 2-3 seedling per hill. The fertilization was applied as recommended rate and time of application. These genotypes have a wide range

of variation due to their different genetic background. Data was recorded on the grain yield (tons/ha), heading date (days), no. productive tillers/plant, plant height(cm), panicle length(cm), panicle weight(g), fertility percentage(%) and 1000 grain weight(g). At first, the data were analyzed by the ordinary analysis of variance to test the significance of differences among the fourteen genotypes. The genotypes mean squares were found to be significant; for all traits studied in their combined data. The combined analysis was calculated over the two years to test the interaction of the different genetic components with the two years as two different environmental conditions. Homogeneity test was done followed by Bartlett (1937) before one proceeds with the computations of the combined experiments and the error variances of the tests were homogenous.

In 2005, seeds of the above mentioned varieties were grown in three successive sowing dates with 10 days intervals to overcome the difference of heading date among them. Thirty days after sowing, seedlings of each variety were individually transplanted in the permanent field in three rows, five meters long and 20 cm between plants and rows. Crossing was carried out between some Delta-Med varieties and Egyptian parents to produce F1 hybrid seeds. Bulk emasculation method was practiced by using hot water technique. The hybrid seeds were grown in the winter glass house to produce F1 generations and the normal F1 at the experimental farm at 2006. The list of the crosses made during 2005 rice growing season are presented in Table 3. In 2006 season, a total of 24 crosses besides their parents were grown in a randomized complete block design (RCBD) with three replications, each replicate consisted of five rows for each parent and F1 cross. Each row was 5 m long and contained 25 hills with 20 cm between rows. Weeds were chemically controlled by applying two liters of Saturn/feddan four days after transplanting. Nitrogen fertilizer was applied as recommended rate and time of application. The crosses listed in Table 3 was designed as to fulfill most of our breeding objectives, *i.e.*, yield potential, resistance to biotic stresses, earliness and grain quality parameters. The hybrid seeds of these crosses were divided in two groups, the first was grown in the winter nursery of the glass house of RRTC to be grown as F2 in the Experimental farm of RRTC in 2006. The second group was kept to be grown as F1 generation in 2006. The F1 and F2 crosses were subjected to pedigree method for selection in the segregating generations. This allows additive alleles to be accumulated in such complex traits. The F4, F5 and F6 generations in 2010 which have been grown as F5, F6 and F7 in 2011 rice growing season. The data of each character of each F7 hybrid population (individuals) were analyzed separately by the analysis of variance, according to Panse & Sukhatme (1957). The genetic parameters, *i.e.*, phenotypic variance, genotypic variance, genetic coefficient of variation was computed according to the formula suggested by Burton (1952). Heritability in broad sense and genetic advance upon selection were computed according to Johanson *et al.* (1955).

TABLE 3. List of crosses made during 2005 season between Delta -Med and Egyptian rice varieties .

No.	Crosses	Source
1	Bahia / Giza 177	NC – 2005 – 51
2	Bahia / Giza 178	NC – 2005 – 79
3	Bahia / Sakha 101	NC – 2005 – 126
4	Bahia / Sakha 102	NC – 2005 – 147
5	Bahia / Sakha 103	NC – 2005 – 172
6	Bahia / Sakha 104	NC – 2005 – 196
7	Bahia / Egyptian Yasmine	NC – 2005 – 200
8	Bahia / Yun Len 18	NC – 2005 – 229
9	Bahia / Geyhwa 7	NC – 2005 – 230
10	Bahia / Onpo 1	NC – 2005 – 231
11	Bahia / GZ 6522-15-1-1-3	NC – 2005 – 277
12	Bahia / GZ 7842-4-1-3-1	NC – 2005 – 317
13	Bahia / CT 9852-B-2-1-2-4PM	NC – 2005 – 359
14	Bomba / Giza 177	NC – 2005 – 52
15	Bomba / GZ 1368-S-5-4	NC – 2005 – 100
16	Bomba / Sakha 101	NC – 2005 – 127
17	Bomba / Sakha 102	NC – 2005 – 148
18	Bomba / Sakha 103	NC – 2005 – 173
19	Bomba / Sakha 104	NC – 2005 – 197
20	Bomba / Balado	NC – 2005 – 228
21	Bomba / GZ 6522-15-1-1-3	NC – 2005 – 278
22	Bomba / GZ 7842-4-1-3-1	NC – 2005 – 318
23	Bomba / GZ 7456-13-6-5-3-	NC – 2005 – 335
24	Bomba / GZ 7563-20-10-8-1	NC – 2005 – 381

Results and Discussion

Characters of Delta-Med materials grown

Results of the evaluation experiment are presented in Table 4. For the 3rd year of evaluation, the Egyptian varieties exceeded the Delta-Med entries in grain yield and its components except 1000 grain weight. All Egyptian varieties exceeded 10 tons / ha. Giza 181 and Sakha 101 recorded the highest grain yield (12.43 and 12.30 ton/ ha, respectively). For Spanish and Delta med genotypes, grain yield were generally low and ranged between 6.57 tons for Tebre and 9.54 tons /ha for Publa. The Delta- Med genotypes had generally shorter duration and higher 1000 grain weight (Table 4). 1000 grain weight (g) for Delta med genotypes ranged from 20.70 in Puntal to 36.10 in 80116-TR259-1-1, while in case of Egyptian rice varieties, 1000 grain weight values ranged from 21.90 in Giza 178 to 28.80 in Sakha 101.

This variation detected among the Egyptian and Delta-Med genotypes will give a great opportunity to enhance the trait value for most of the studied traits by using conventional breeding to utilize such high level of variation among those genotypes.

TABLE 4. Mean performance data of Delta-Med materials grown at Sakha (2010).

Genotype	D.H (day)	P.H (cm)	P.L (cm)	No.till./ plant	G.y (ton/ha)	Fert. (%)	1000-g-w (g)
Giza 177	96.00	100.00	26.00	20.33	9.98	89.10	22.30
Giza178	98.00	102.00	24.60	28.66	10.70	94.00	21.90
Giza182	106.00	98.00	23.60	21.66	9.15	87.92	23.80
Giza 181	112.30	110.00	27.60	25.00	9.52	85.12	26.40
Sakha 101	111.00	91.00	19.60	29.66	11.00	93.95	28.80
Sakha102	98.00	107.00	28.60	26.00	9.50	95.52	27.40
Sakha103	94.00	104.00	26.00	22.66	9.90	86.19	22.10
Sakha104	106.00	108.00	26.30	27.33	10.50	93.64	27.60
Bala	94.00	95.00	22.00	25.00	9.31	95.54	30.40
Susan	94.00	74.60	21.60	23.33	7.79	96.81	27.50
Bahia	91.00	101.00	26.30	13.00	5.10	75.70	32.30
Bomba	92.00	128.30	25.30	20.33	7.79	87.19	27.80
Montsianejj	92.00	78.30	26.00	27.66	5.19	69.81	31.70
Senia	93.00	82.00	21.30	17.33	5.30	86.10	27.60
Tebre	84.00	77.00	21.00	19.66	6.98	87.50	30.50
79004- TR4-4-2-1-1	92.00	111.00	27.00	25.33	8.09	89.03	31.10
79061- TR64-3-3-1-1	89.00	113.00	19.60	18.33	5.69	78.74	35.70
80110- TR 4-1-1	115.00	110.00	22.00	13.00	4.58	71.11	34.20
80116- TR 259-1-1	92.00	99.30	20.00	16.00	5.53	78.65	36.10
96066- TR 1805-5-1-1	95.00	102.60	18.60	10.00	2.88	67.09	30.10
Publa	96.00	98.00	27.30	11.66	3.91	67.16	23.30
Hisprgran	92.00	106.00	19.30	18.30	4.11	88.88	32.80
Thaibonet	98.00	90.60	21.30	22.00	6.00	80.42	31.00
Puntal	103.00	95.00	25.00	25.00	7.47	90.90	20.70
LSD at 0.05	2.50	2.90	2.30	2.55	0.70	5.00	2.20

DH= Days to heading, P.H= Plant height, P.L= Panicle length, No.till.=No. productive tillers/plant, G.y= Grain yield(ton/ha), Fert.= Fertility % and 1000-g-w= 1000 grain weight. The same as below.

Biotic stress reaction of the Delta-Med tested materials

As far as biotic stresses is concerned, Egyptian varieties exhibit also considerably higher level of resistance. The reaction of the tested materials to both blast and stem borer are presented in Table 5. Egyptian rice varieties showed higher level of resistance to blast at both leaf and neck infection except Sakha 101 and Sakha 104. These two varieties started to show a blast resistance breakdown during the last three years. Most of the Delta Med showed high level of susceptibility which may limit their use under such ecosystem.

TABLE 5. Biotic stress reaction of the Delta-Med tested materials under Egyptian conditions (2010).

No.	Genotype	Blast Reaction (leaf)	Blast Reaction (nick)	Stem Borer Reaction
1	Giza 177	1	1	R
2	Giza 178	1	1	MR
3	Giza 181	1	1	R
4	Giza182	1	1	R
5	Sakha 101	4	4	S
6	Sakha 102	1	1	R
7	Sakha 103	1	1	R
8	Sakha 104	5	4	S
9	Bala	7	7	HS
10	Susan	7	7	S
11	Bahia	7	7	HS
12	Bomba	7	7	S
13	Montsianeji	7	7	HS
14	Senia	7	7	HS
15	Tebre	5	7	HS
16	79004-TR4-4-2-1-1	5	7	S
17	79061-TR64-3-3-1-1	7	7	S
18	80110- TR 4-1-1	7	7	S
19	80116- TR 259-1-1	5	5	HS
20	96066- TR1805-5-1-1	5	5	S
21	Publa	7	7	S
22	Hisprgran	5	7	S
23	Thaibonet	7	7	HS
24	Puntal	7	7	S

R= Resistant, MR= Moderate resistant , S= Susceptible and HS= Highly susceptible

Characters of the best selected lines (15 lines) from F6 generation

Data from Table 6 shows that most of the selected lines have desirable values for various traits. The red color lines were superior for most of the studied traits especially for yield and its components .These lines exceeded their parental lines. In addition to its higher yield they were earlier in heading and growth duration around 130 days, heavier in grain weight, high fertility percentage. So, we have succeeded to combine Egyptian and Delta-Med desirable traits in one line. These were promoted to be grown as F7 generation in 2011 rice growing season under normal conditions besides the Egyptian rice varieties for comparison. The Selection has been made for yield and its components as well as blast and stem borer resistance.

TABLE 6. The best selected lines (15 lines) from F6 generation (2010).

No.	Crosses	D.H (day)	P.H (cm)	P.L (cm)	No.till/ plant	G.y (ton/ha)	Fert. (%)	1000-g-w (g)	
7	Bomba x Giza182	99.00	93.00	20.00	21.00	7.90	87.00	23.00	
18		102.00	91.00	19.00	25.00	10.58	88.00	25.00	
21		100.00	102.00	20.00	22.00	10.45	90.00	26.00	
25		100.00	100.00	19.00	19.00	8.89	91.00	26.00	
26		99.00	95.00	21.00	26.00	10.87	93.00	26.00	
27		99.00	110.00	23.00	19.00	9.19	89.00	25.00	
28		100.00	88.00	22.00	20.00	8.02	86.00	25.00	
29		101.00	97.00	27.00	30.00	11.22	92.00	27.00	
30		100.00	105.00	25.00	29.00	11.38	94.00	28.00	
39		100.00	100.00	26.00	22.00	8.49	87.00	26.00	
44		100.00	90.00	28.00	28.00	11.00	92.00	29.00	
53		Bomba x IR6554-22- 2-3	100.00	93.00	25.00	24.00	10.75	94.00	27.00
54			99.00	98.00	24.00	31.00	11.14	91.00	28.00
56	98.00		105.00	23.00	30.00	11.03	90.00	27.00	
65	Susan x Giza182	99.00	100.00	22.00	18.00	7.90	88.00	25.00	

Phenotypic correlation coefficients

Phenotypic correlation coefficients among the traits studied are shown in Table 7. Days to heading is significantly and highly significantly and positively associated with no. panicles/plant and fertility %. Plant height was positively and significantly associated with panicle length. Panicle length was highly significantly and positively correlated with grain yield per plant. No. panicles/plant was positively correlated with grain yield per plant and fertility %. One thousand grain weight and fertility percentage were significant and positively correlated with grain yield per plant. These results were in agreement with those reported by Bhattacharyya & Ghosh (2004), Shashidhar *et al.* (2005), Surek & Beser (2003) and Yogameenakshi *et al.* (2004).

TABLE 7. Correlation coefficients among the traits studied of the selected lines from F7 generation.

Characters	D.H	P.H.	P. L	No.panicle/ plant	Fert.	1000- g-w	Grain yield /plant
D.H.	1.000	0.230	0.310	0.520**	0.220*	0.180	0.300
P.H.		1.000	0.780**	0.110	-0.220	0.250	0.280
P.L.			1.000	0.150	0.310	0.230	0.550*
No.panicles /plant				1.000	0.475*	0.210	0.780**
Fert.					1.000	0.225	0.650**
1000-g-w						1.000	0.850**
Grainyield/plant							1.000

*Significant and ** highly significant

Genetic components

In general, the phenotypic coefficient of variation (PCV), (Table 8) was high for fertility % (110.00), number of panicles per plant (100.00), and plant height (98.00), while it was low for panicle length (40.00), days to heading (50.00) and grain yield/plant (55.00). Genotypic coefficient of variation GCV showed the same trend but the corresponding values were lower thereby implying the influence of environment on the genotypes. Heritability estimates, in general, were moderate to high for all characters studied, the values ranged from 62.00% to 76.53%. Reasonable values of genetic advance were found for all traits studied; implying that higher proportion of genotypic variance was attributed to additive genetic variance, and therefore, selection would be effective and useful for these traits from the progenies which will be produced. Sheeba (2005) and Raju *et al.* (2004) reported the same results with the other varieties.

TABLE 8. Genetic parameters of variation for some characters studied in the promising lines.

Characters	Genotypic variance (%)	Phenotypic variance(%)	Heritability in broad sense(%)	Genetic advance
Days to heading(day)	35.00	50.00	70.00	10.19
Plant height(cm)	75.00	98.00	76.53	15.50
Panicle length(cm)	25.00	40.00	62.50	8.07
No. panicles/plant	68.10	100.00	68.00	14.00
100 grain weight (gram)	66.00	86.00	76.74	14.51
Fertility (%)	85.00	110.00	77.27	16.63
Grain yield(ton/ha)	38.00	55.00	69.00	10.54

The best five lines are presented in Table 9. These lines can be distributed at the different rice growing areas in Egypt as well as can be tested at the Delta-Med conditions to be used as a donor parents in rice breeding program or as new rice varieties directly possess high yielding ability, good grain quality characters and blast and stem borer resistance.

TABLE 9. The best selected lines (5 lines) from F7 generation (2011).

No.	Crosses	D.H (day)	P.H (cm)	P.L (cm)	No.till/ plant	G.y (ton/ha)	Fert. (%)	1000-g-w (g)
26	Bomba x Giza182	100.00	98.00	22.00	24.00	10.71	90.00	26.00
29		103.00	100.00	25.00	27.00	11.18	90.00	27.00
30		98.00	107.00	23.00	28.00	11.00	91.00	28.00
44		100.00	93.00	26.00	25.00	10.60	92.00	29.00
53	Bomba x	102.00	90.00	25.00	22.00	10.50	92.00	27.00
56	IR6554-22-2-3	95.00	100.00	23.00	28.00	11.00	90.00	27.00

Conclusion

Data obtained from this study show that most of the selected lines have desirable values for various traits. Some lines were superior for most of the studied traits especially for yield and its components. These lines exceeded their

parental lines. In addition to its higher yield they were earlier in heading and growth duration around 130 days, heavier in grain weight, high fertility percentage. So, we have succeeded to combine Egyptian and Delta-Med desirable traits in one line. Now we have promising lines from the advanced population's possess many desirable traits. Results of this cooperation will support development in the Mediterranean area and Delta Med areas. We are going to select the most promising lines from the advanced generation to be grown under different locations in some Delta Med countries to test their performance for grain yield, stem borer and blast resistance.

References

- Bhattacharyya, S. and Ghosh, S.K. (2004)** Association among yield related traits of twenty four diverse land races of rice. *Crop Res.* **27**, 90-93.
- Bartlett, M.S. (1937)** Properties of sufficiency and statistical tests. *Proc. Roy. Soc., London Series A* **160**, pp. 268-282.
- Boonjung, H. and Fukai, S. (1996)** Effect of soil water deficit at different growth stage on rice growth and yield under upland conditions. *Field Crops Res.* **48**ss, pp.47-55.
- Burton, G.W. and Devane, E.H. (1952)** Estimating heritability in tall fescue from replicated clonal material. *Agron. J.* **45**, 478-841.
- International Rice Research Institute (1996)** Standard Evaluation System of Rice
- Jacob, W. Kijne (2006)** A biotic stress and water scarcity: Identifying and resolving conflicts from plant level to global level. *Field Crops Research* **97**, 3-18.
- Johanson, H.W., Robinson, H.F. and Comstock, R.F. (1955)** Estimates of genetic and environmental variability in Soybeans. *Agron. J.* **47**, 314-318.
- Khunthasuvon, S., Rajatasereekul, S., Hanviriyapant, P. and Basnayake, J. (1998)** Effects of fertilizer application and irrigation on grain yield of several rice cultivars. *Field Crops Res.* **59**, 99-108.
- Panse and Sukhatme (1957)** "Statistical Methods for Agriculture Workers". India Council of Agric. Res. New Delhi, India.
- Raju, C.H.S., Rao, M.V.B. and Sundarshanam, A. (2004)** Genetic analysis and character association in F₂ generation of rice. *Madras Agric. J.* **91**, 66-69.
- RRTC (2007)** The 11th Annual Workshop of the National Rice Research Program, Rice Research and Training Center, Sakha, Egypt, Jan., 22nd, 2008.
- Shashidhar, H. E., Pasha, F., Manjunath, Janamatti, M. S. Vinod and Adnan Kanbar (2005)** Correlation and path coefficient analysis in traditional cultivars and doubled haploid lines of rainfed lowland rice (*Oryza sativa* L.). *Oryza*, **42**, 156-159.

- Sheeba, A. (2005)** Genetic studies on drought tolerance and stability of Temperature Sensitive Genic Male Sterility (TGMS) based rice hybrids. *Ph.D. Thesis*, TNAU, Coimbatore.
- Surek, H. and Beser, N. (2003)** Correlation and path analysis for some yield related traits under rainfed condition. *Turkish J. Agric. Forestry*, **27**, 77-83.
- Yogameenakshi, P., Nadarajan, N. and Anbumalarmathi, J. (2004)** Correlation and path analysis on yield and drought tolerant attributes in rice (*Oryza sativa* L.). under drought stress. *Oryza*, **41**, 68-70.

(Received 3/10/2012;
accepted 20/2/2013)

استراتيجيات ومنهجيات تحسين أصناف الأرز في مصر

عبد الله عبد النبي عبد الله، شيماء عبد العظيم بدوي* ، عبد العظيم الطنطاوي بدوي
قسم بحوث الأرز - معهد بحوث المحاصيل - مركز البحوث الزراعية - الجيزة
وقسم المحاصيل - كلية الزراعة - جامعة كفر الشيخ - مصر .

أجريت هذه الدراسة في مزرعة مركز البحوث والتدريب في الأرز - سخا خلال
موسم زراعة الأرز من ٢٠٠٥ إلى ٢٠١١. وكان الهدف من الدراسة هو تقييم
مجموعة من الأصناف المنزوعة في دول حوض البحر الأبيض المتوسط (الدلتا
ميد) تحت الظروف المصرية وأيضاً استنباط سلالات جديدة من الأرز عن طريق
التهجين بين تلك الأصناف والأصناف المصرية المحلية.

أوضحت النتائج وجود اختلافات معنوية بين الأصناف المصرية واصناف
الدلتا ميد وما بين اصناف كل مجموعة .

تفوقت الأصناف المصرية على اصناف الدلتاميد في صفة المحصول
ومكوناته ، اما اصناف الدلتا ميد فكانت بصفة عامة مبكرة النضج ومنخفضة في
المحصول وذات وزن عالي للحبوب.

وبالتهجين بين تلك الأصناف تم التوصل إلى سلالات مبشرة تجمع ما بين
الصفات المرغوبة في كلا المجموعتين ، وتم تقييم الانسال الناتجة من الهجن
بطريقة التربية باستخدام سجلات النسب تحت الظروف العادية .

تم انتخاب احسن السلالات من الجيل السادس في ٢٠١٠ لتزرع كنباتات جيل
سابع في موسم ٢٠١١. أوضحت النتائج أن تلك السلالات كانت مقاومة لمرض
الفحة بالإضافة إلى المحصول العالي ، ووجد أيضاً قيم عالية للتباين المظهري
والتباين الوراثي ودرجة التوريث لكل الصفات المدروسة .